

Insolvency - (Quasi-)Poisson Model and Negative Binomial Model

February 5, 2020

First the insolvency data are loaded:

```
> library(catdata)
> data(insolvency)
> attach(insolvency)
```

For the number of insolvent firms between 1994 and 1996 a Poisson model is fitted with time as predictor. Time is considered as a number from 1 to 36, denoting the month from January 1994 to December 1996.

```
> ins1 <- glm(insolv ~ case + I(case^2), family=poisson(link=log), data=insolvency)
> summary(ins1)

Call:
glm(formula = insolv ~ case + I(case^2), family = poisson(link = log),
     data = insolvency)

Deviance Residuals:
    Min      1Q      Median      3Q      Max 
-3.2037 -0.9083 -0.2517  0.4880  3.0340 

Coefficients:
            Estimate Std. Error z value Pr(>|z|)    
(Intercept) 4.1916952  0.0617994 67.827 < 2e-16 ***
case        0.0197825  0.0073901  2.677  0.00743 **  
I(case^2)   -0.0002670  0.0001896 -1.409  0.15897  
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for poisson family taken to be 1)


Null deviance: 108.128  on 35  degrees of freedom
Residual deviance: 75.287  on 33  degrees of freedom
AIC: 306.82

Number of Fisher Scoring iterations: 4

> # plot(ins1)
```

Scatter-Plot of number of insolvent firms dependent of the month (1-36). With estimated curve of the log-linear model.

```
> plot(case, insolv)
> points(ins1$fitted.values, type="l")
```



In many real-world datasets the variance of count-data is higher than predicted by the Poisson distribution. So next a Poisson model with dispersion parameter is fitted (Quasi-Poisson model).

```
> ins2 <- glm(insolv ~ case + I(case^2), family=quasipoisson, data=insolvency)
> summary(ins2)

Call:
glm(formula = insolv ~ case + I(case^2), family = quasipoisson,
     data = insolvency)

Deviance Residuals:
    Min      1Q      Median      3Q      Max 
-3.2037 -0.9083 -0.2517  0.4880  3.0340 

Coefficients:
            Estimate Std. Error t value Pr(>|t|)    
(Intercept) 4.1916952  0.0939826 44.601   <2e-16 ***
case        0.0197825  0.0112387  1.760   0.0876 .  
I(case^2)  -0.0002670  0.0002883 -0.926   0.3611  
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
(Dispersion parameter for quasipoisson family taken to be 2.312738)
```

```
Null deviance: 108.128 on 35 degrees of freedom
Residual deviance: 75.287 on 33 degrees of freedom
AIC: NA
```

```
Number of Fisher Scoring iterations: 4
```

```
> # plot(ins2)
```

An alternative to a quasi-poisson model is to use the negative binomial distribution.

```
> library(MASS)
> ins3 <- glm.nb(insolv ~ case + I(case^2), data=insolvency)
> summary(ins3)
```

Call:

```
glm.nb(formula = insolv ~ case + I(case^2), data = insolvency,
       init.theta = 77.92952593, link = log)
```

Deviance Residuals:

Min	1Q	Median	3Q	Max
-2.3666	-0.6333	-0.1710	0.3350	2.0042

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	4.1953863	0.0861256	48.712	<2e-16 ***
case	0.0192833	0.0105170	1.834	0.0667 .
I(case^2)	-0.0002546	0.0002728	-0.933	0.3506

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```
(Dispersion parameter for Negative Binomial(77.9295) family taken to be 1)
```

```
Null deviance: 52.104 on 35 degrees of freedom
Residual deviance: 36.312 on 33 degrees of freedom
AIC: 296.27
```

```
Number of Fisher Scoring iterations: 1
```

```
Theta: 77.9
Std. Err.: 35.5
```

2 x log-likelihood: -288.269

Since counts are rather large in addition a normal distribution model is fitted.

```
> ins4 <- glm(insolv ~ case + I(case^2), family=gaussian(link=log), data=insolvency)
> summary(ins4)
```

```

Call:
glm(formula = insolv ~ case + I(case^2), family = gaussian(link = log),
     data = insolvency)

Deviance Residuals:
    Min      1Q  Median      3Q      Max
-25.809   -8.744   -2.374    4.560   30.480

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) 4.1836089  0.1005663  41.600 <2e-16 ***
case        0.0208026  0.0115423   1.802  0.0806 .
I(case^2)  -0.0002915  0.0002896  -1.007  0.3214
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for gaussian family taken to be 193.2793)

Null deviance: 9147.0  on 35  degrees of freedom
Residual deviance: 6378.1  on 33  degrees of freedom
AIC: 296.54

Number of Fisher Scoring iterations: 4

```